

LINEARIZATION OF PROPER GROUPOIDS

NGUYEN TIEN ZUNG

ABSTRACT. We prove the following result, conjectured by Alan Weinstein: every smooth proper Lie groupoid near a fixed point is locally linearizable, i.e. it is locally isomorphic to the associated groupoid of a linear action of a compact Lie group. In combination with a slice theorem of Weinstein, our result implies the smooth linearizability of a proper Lie groupoid in the neighborhood of an orbit under a mild condition.

1. INTRODUCTION

In the theory of Lie groupoids, the role played by proper Lie groupoids is probably comparable to that played by compact Lie groups in the Lie group theory. However, while compact Lie groups have been classified completely long time ago, the study of proper Lie groupoids has been initiated only recently by Alan Weinstein [9, 11], and many basic questions remained open. One of them, formulated by Weinstein as the *proper groupoid structure conjecture*, is about the local linearizability of proper Lie groupoids near fixed points. The main purpose of the present paper is to prove this conjecture.

So let us consider a smooth proper groupoid near a fixed point. It means that we are given the following data:

i) A smooth manifold M (the *total space*), a smooth submanifold B of M (the *base space*, also called the *space of units*), and two smooth maps $M \xrightarrow{s} B$, $M \xrightarrow{t} B$, called the *source map* and the *target map* respectively. The restrictions of s and t on B are identity. Here, by *smooth* we mean either C^∞ -smooth, or C^h -smooth for some natural number h .

ii) Denote by M_k ($k \geq 2$) the subset of $M \times \cdots \times M$ (k times) consisting of k -tuples $(p_1, \dots, p_k)$ of elements of M such that $s(p_1) = t(p_2)$, $s(p_2) = t(p_3)$, $\dots$, $s(p_{k-1}) = t(p_k)$. Then there is a map from M_2 to M , called the *product map*: if $(p, q) \in M_2$, i.e. $s(p) = t(q)$, then we can form the product of p with q , denoted by $p.q$.

iii) There is a map from M to itself, called the *inversion*, which associates to each element $p \in M$ an element denoted by p^{-1} , called its inverse. The inverse of each point in B is itself.

Date: Second version, May/2003.

1991 Mathematics Subject Classification. 58H05, 57S15.

Key words and phrases. compact Lie groups, proper groupoids, linearization, averaging, near homomorphisms.

The product map and the inversion map satisfy the following conditions: $s(p.q) = s(q)$ and $t(p.q) = t(p)$ for any $(p, q) \in M_2$; $(p.q).r = p.(q.r)$ for any $(p, q, r) \in M_3$; $p.p^{-1} \in B$ and $p^{-1}.p \in B$ for any $p \in M$; $p.q = p$ if $q \in B$ and $p.q = q$ if $q \in B$.

iv) A point $x_0 \in B$, which is a fixed point for the groupoid, in the sense that for any $p \in M$ with $s(p) = x_0$ we also have $t(p) = x_0$. The set $G = \{p \in M, s(p) = x_0\}$ is then a group under the above product map.

The properness conditions that we impose on G and s and t are that G is a compact (not necessarily connected) Lie group, and $M \xrightarrow{s} B$, $M \xrightarrow{t} B$ are smooth fiber bundles, with fibers diffeomorphic to $G = s^{-1}(x_0) = t^{-1}(x_0)$.

For each point $x \in B$, the *orbit* of our groupoid via x is the set $O(x) := t(s^{-1}(x))$, which is the same as $s(t^{-1}(x))$. A subset in B is called *invariant* if it is saturated by the orbits of the groupoid. As observed by Weinstein [11], any neighborhood of x_0 in B contains an invariant neighborhood (simply take a sufficiently small open ball $D \subset B$ which contains x_0 and take $D' = t(s^{-1}(D))$, then $D' \supset D$ is a small invariant open neighborhood of x_0). Of course, the restriction of a groupoid to an invariant set is still a groupoid. Since we are interested in the local behavior of our proper groupoid $M \rightrightarrows B$ near $G = s^{-1}(x_0)$, we can assume that B is a small neighborhood of x_0 in an Euclidean space.

If the compact Lie group G acts smoothly on a manifold X , then we can define the associated *action groupoid* $G \times X \rightrightarrows X$: the source map is $s(g, x) = x$ for $(g, x) \in G \times X$, the target map is $t(g, x) = g.x$ (the action of g on x), and the inversion map is $(g, x)^{-1} = (g^{-1}, g.x)$. The space of units is $\{1_G\} \times X$ (where 1_G denotes the neutral element of G), which is identified with X . If the action of G on X has a fixed point, then due to Bochner's linearization theorem [2], we may assume that X is a small neighborhood of this fixed point in an Euclidean space, and the action of G on X is linear. In this case we say that the action groupoid $G \times X \rightrightarrows X$ is a *linear action groupoid*. We will say that our proper groupoid $M \rightrightarrows B$ is isomorphic to an action groupoid $G \times X \rightrightarrows X$ if there is a smooth diffeomorphism from M to $G \times X$ which preserves the maps s, t , the product and the inversion.

The main result of the present paper is the following

Theorem 1.1. *Let $M \rightrightarrows B$ be a C^h (resp., C^∞) smooth proper groupoid with a fixed point $x_0 \in B$, with the notations as of above. Then the restriction of this groupoid to a sufficiently small invariant neighborhood of x_0 in B is C^h (resp., C^∞) isomorphic to a linear action groupoid of G .*

An important application of Theorem 1.1 is in the study of symplectic groupoids, where one can show that proper symplectic groupoids are locally isomorphic to standard models, see [14, 13]. In turn, this result about the local structure of proper symplectic groupoids is important for the study of intrinsic convexity properties of momentum maps, initiated by Weinstein [9, 10], see [14, 13].

Theorem 1.1, together with Weinstein's slice theorem (Theorem 9.1 of [11]), immediately implies the following result about the local linearization of a proper groupoid in the neighborhood of an orbit:

Theorem 1.2. *Let $\Gamma \rightrightarrows X$ be a proper Lie groupoid, and let $\mathcal{O}$ be an orbit of Γ which is a manifold of finite type. Then there is a neighborhood $\mathcal{U}$ of $\mathcal{O}$ in X such that the restriction of Γ to $\mathcal{U}$ is isomorphic to the restriction of the action groupoid $\Gamma_{\mathcal{O}} \times_{\mathcal{O}} N\mathcal{O} \rightrightarrows N\mathcal{O}$ to a neighborhood of the zero section.*

In Theorem 1.2, $N\mathcal{O}$ means the normal bundle of $\mathcal{O}$ in X , and $\Gamma_{\mathcal{O}}$ is the restriction of Γ to $\mathcal{O}$. The condition that $\mathcal{O}$ is of finite type means that there is a proper map from $\mathcal{O}$ to $\mathbb{R}$ with a finite number of critical points. See [11] for details.

Theorem 1.1 is essentially equivalent to the existence of a smooth *surjective homomorphism* ϕ from M to G (after shrinking B to a sufficiently small invariant neighborhood of x_0), i.e. a smooth map $\phi : M \rightarrow G$ which satisfies

$$(1.1) \quad \phi(p.q) = \phi(p).\phi(q) \quad \forall (p, q) \in M_2 ,$$

such that the restriction of ϕ to $G = s^{-1}(x_0) \subset M$ is an automorphism of G . We may assume that this automorphism is identity.

Indeed, if there is an isomorphism from M to an action groupoid with the total space $G \times X$, then the composition of the isomorphism map $M \rightarrow G \times X$ with the projection $G \times X \rightarrow G$ is such a homomorphism. Conversely, if we have a homomorphism $\phi : M \rightarrow G$, whose restriction to $G = s^{-1}(x_0) \subset M$ is the identity map of G , then assuming that B is a sufficiently small invariant neighborhood of x_0 , we have a diffeomorphism

$$(1.2) \quad (\phi, s) : M \rightarrow G \times B .$$

Denote by θ the inverse map of (ϕ, s) . Then there is an action of G on B defined by $g.x = t(\theta(g, x))$, and the map (ϕ, s) will be an isomorphism from $M \rightrightarrows B$ to the action groupoid $G \times B \rightrightarrows B$. This action groupoid is linearizable by Bochner's theorem, implying that the groupoid $M \rightrightarrows B$ is linearizable.

In order to find such a homomorphism from M to G , we will use the averaging method. The idea is to start from an arbitrary smooth map $\phi : M \rightarrow G$ such that $\phi|_G = Id$ (Recall that we identify $s^{-1}(x_0) = t^{-1}(x_0)$ with G). Then Equality (1.1) is not satisfied in general, but it is satisfied for $p, q \in G$. Hence it is "nearly satisfied" in a small neighborhood of $G = s^{-1}(x_0)$ in M . In other words, if the base B is small enough, then $\phi(p.q)\phi(q)^{-1}$ is near $\phi(p)$ for any $(p, q) \in M_2$. We will replace $\phi(p)$ by the average value of $\phi(p.q)\phi(q)^{-1}$ for q running on $t^{-1}(s(p))$ (it is to be made precise how to define this average value). This way we obtain a new map $\hat{\phi} : M \rightarrow G$, which will be shown to be "closer" to a homomorphism than

the original map ϕ . By iterating the process and taking the limit, we will obtain a true homomorphism ϕ_∞ from M to G .

Let us mention that the averaging method is a classical method which is relatively simple and effective in many problems involving compact group actions. For example, it was used by Elie Cartan [3, 4] to prove the existence of a fixed point of a compact group action under some conditions (see [8] and references therein for recent results on invariant manifolds of compact group actions). Bochner's linearization theorem [2] is indeed based on a very simple averaging formula. The iterative averaging method for was already used by Grove, Karcher and Ruh in [6] to show that near-homomorphisms between compact Lie groups can be approximated by homomorphisms. Weinstein [8, 9, 11] also suggested the averaging method for proving Theorem 1.1, although he considered near-homomorphisms from G to the group of bisections of $M \rightrightarrows B$ instead of near-homomorphisms from M to G . In this paper, instead of using the Haar measure of a compact group for making the averaging, we will use a "Haar system" on a proper groupoid. More precisely, we will use a family of probability measures on the fibers of the fibration $M \xrightarrow{t} B$, which are invariant under left translations. This "technical" detail allows us to effectively control the convergence of our iterative averaging process, using standard Banach-space estimations.

It is clear (see, e.g., [9]) that the problem of linearization of Lie groupoids is closely related to the problem of linearization of Lie algebroids. In this aspect, Theorem 1.1 is related to our results with Philippe Monnier [12, 7] on Levi decomposition of Lie algebroids. Several natural questions arise, among which: is there a similar result about Levi decomposition of Lie groupoids near a fixed point? If it exists, can we prove it by the averaging method? Let us mention that the results of [12, 7] are obtained with the aid of the fast convergence methods of Kolmogorov and Nash-Moser, which are technically considerably more complicated than the averaging method. Theorem 1.1 or a more general result about Levi decomposition of Lie groupoids, together with criteria about (partial) integrability of Lie algebroids like the one of Crainic-Fernandes [5], might eventually lead to a simplification and improvement of our papers [12, 7].

The rest of this paper is devoted to the construction of a smooth homomorphism from M to G (after shrinking the base B if necessary), whose restriction to G is the identity map of G . As explained above, Theorem 1.1 follows immediately from the existence of such a homomorphism. In Section 2 we present our iterative averaging algorithm, and in Section 3 we use standard analytical estimates to show that our algorithm actually yields a (C^h or C^∞) smooth homomorphism.

In this paper, we will consider only C^h and C^∞ smooth groupoids, but we suspect that, with more care, one can probably show that our algorithm for constructing a linearization of $M \rightrightarrows B$ works in the real-analytic category as well.

2. THE AVERAGING ALGORITHM

2.1. A translation-invariant measure. For each point $p \in M$ denote by $T(p) = t^{-1}(t(p))$ (resp., $S(p) = s^{-1}(s(p))$) the fiber of the fibration $M \xrightarrow{t} B$ (resp. $M \xrightarrow{s} B$) which contains p .

Each element $q \in M$ defines a *left translation map* τ_q from $T(s(q))$ to $T(q)$ as follows: $\tau_q(r) = q.r$ for $r \in T(s(q))$.

Let us fix on M an arbitrary smooth Riemannian metric. (Throughout this paper, smooth objects have the same smoothness class as that of our groupoid $M \rightrightarrows P$, unless mentioned explicitly otherwise). It induces on each fiber $T(p)$ of the fibration $M \xrightarrow{t} B$ a smooth measure (which depends smoothly on the fiber itself), which we will denote by $d\mu_0$. Similarly, the metric on M induces on each fiber $S(p)$ of the fibration $M \xrightarrow{s} B$ a smooth measure, which we will denote by $d\nu_0$.

Lemma 2.1. *There is a smooth positive function f on M such that the measure $d\mu = fd\mu_0$ defined on each fiber of the fibration $M \xrightarrow{t} B$ is a translation-invariant probability measure: for each $q \in M$ the left translation map τ_q from $T(s(q))$ to $T(q)$ preserves the measure $d\mu$, and moreover $\int_{T(q)} 1d\mu = 1$.*

Translation-invariant measures are also called *Haar systems* (see, e.g., [1]), and the above lemma is probably a well-known folklore result, but to make our paper self-contained, we will include a simple proof of it here.

Proof. Define $\tilde{f}$ by the following integral formula:

$$(2.1) \quad \tilde{f}(r) = \int_{p \in S(r)} \frac{\tau_{p.r^{-1}}^* d\mu_0(r)}{d\mu_0(r)} d\nu_0 .$$

In the above formula, $\tau_{p.r^{-1}}$ is the left translation map from $T(r)$ to $T(p)$, $\tau_{p.r^{-1}}^* d\mu_0(r)$ means the pull-back at r of the smooth measure $d\mu_0$ on $T(p)$ by $\tau_{p.r^{-1}}$, and $d\nu_0$ is the measure on $S(r)$ induced by the metric on M . One checks immediately that $\tilde{f}$ is a smooth positive function on M , and the measure $d\tilde{\mu} = \tilde{f}d\mu_0$ on the fibers of $M \xrightarrow{t} B$ is translation invariant. Dividing $\tilde{f}$ by the integral $I = \int_{T(p)} \tilde{f}d\mu_0$, we get the required smooth function $f = \tilde{f}/I$ and translation-invariant measure $d\mu = fd\mu_0$. $\square$

From now on, we will fix a measure $d\mu$ provided by the above lemma. We will use it in our averaging formulas. The invariance of $d\mu$ under translations will be important for us.

Remark 2.2. As pointed out to me by A. Weinstein, there is a natural 1-1 correspondence between smooth translation-invariant volume forms on fibers of $M \xrightarrow{t} B$ and smooth non-vanishing sections of the top exterior power of the associated Lie algebroid over B . This point of view gives another simple

proof of the above lemma, without the need of choosing a Riemannian metric on M .

Remark 2.3. Another simple way to define a measure on $T(q)$ is to restrict the map $\phi : M \rightarrow G$ to $T(q)$ and take the pull-back of the Haar measure of G via this restricted map. But this measure is not translation-invariant, and moreover it depends on the differential of ϕ . Maybe the averaging algorithm presented in the next subsection still works with this induced measure from G , but the estimates involved will be much more complicated, and one might have to use Nash-Moser theory due to “loss of differentiability” in some inequalities. (This loss is hidden in the measure, which depends on the differential of ϕ).

2.2. The averaging formula. We fix a bi-invariant metric on the Lie algebra $\mathfrak{g}$ of G and the induced bi-invariant metric on G itself. Denote by 1_G the neutral element of G . For each number $\rho > 0$, denote by $B_{\mathfrak{g}}(\rho)$ (resp., $B_G(\rho)$) the closed ball of radius ρ in $\mathfrak{g}$ (resp., G) centered at 0 (resp., 1_G). By rescaling the metric if necessary, we will assume that the exponential map

$$(2.2) \quad \exp : B_{\mathfrak{g}}(1) \rightarrow B_G(1)$$

is a diffeomorphism. Denote by

$$(2.3) \quad \log : B_G(1) \rightarrow B_{\mathfrak{g}}(1)$$

the inverse of $\exp$. Define the distance $\Delta(\phi)$ of ϕ from being a homomorphism as follows:

$$(2.4) \quad \Delta(\phi) = \sup_{(p,q) \in M_2} d(\phi(p.q).\phi(q)^{-1}.\phi(p)^{-1}, 1_G) .$$

Let $\phi : M \rightarrow G$ be a smooth map such that $\phi|_G$ is identity. We will assume that $\Delta(\phi) \leq 1$, so that the following map $\hat{\phi} : M \rightarrow G$ is clearly well-defined:

$$(2.5) \quad \hat{\phi}(p) = \exp\left(\int_{q \in T(s(p))} \log(\phi(p.q).\phi(q)^{-1}.\phi(p)^{-1})d\mu\right).\phi(p) .$$

Since $d\mu$ is translation-invariant, by the change of variable $r = p.q$, we can also write $\hat{\phi}$ as:

$$(2.6) \quad \hat{\phi}(p) = \exp\left(\int_{r \in T(p)} \log(\phi(r).\phi(p^{-1}.r)^{-1}.\phi(p)^{-1})d\mu\right).\phi(p) .$$

Due to the commutativity of the maps $\exp$ and $\log$ with the adjoint actions, we can also write $\hat{\phi}$ as:

$$(2.7) \quad \hat{\phi}(p) = \phi(p) \cdot \exp\left(\int_{q \in T(s(p))} \log(\phi(p)^{-1}.\phi(p.q).\phi(q)^{-1})d\mu\right) .$$

It is clear that $\widehat{\phi}$ is a smooth map from M to G , and its restriction to $G = s^{-1}(0) \subset M$ is also identity.

2.3. The iterative process. Starting from an arbitrary ϕ (such that $\phi|_G = Id$) as above, we construct a sequence of maps $\phi_n : M \rightarrow G$, by the recurrence formula $\phi_1 = \phi$, $\phi_{n+1} = \widehat{\phi_n}$. In the next section we will show that this sequence is well-defined (after a restriction of B to a smaller invariant neighborhood of x_0 if necessary), and that

$$(2.8) \quad \phi_\infty = \lim_{n \rightarrow \infty} \phi_n$$

exists, is smooth, and is a homomorphism from M to G .

3. PROOF OF CONVERGENCE

3.1. Spaces of maps and C^k -norms. As in the introduction, for each $n \in \mathbb{N}$, we denote by M_n the space of n -tuples of points $(p_1, \dots, p_n)$ of M such that $s(p_1) = t(p_2), s(p_2) = t(p_3), \dots, s(p_{n-1}) = t(p_n)$. There are many natural product and projection maps among these spaces M_n ($n \in \mathbb{N}$), and these maps are smooth. The reader may recall that these spaces M_n , together with natural maps among them, are used in the definition of the cohomology of M , though we will not use any cohomology (at least not in an explicit way) here. For each $n \in \mathbb{N}$, the manifold M_n is smoothly diffeomorphic to $B \times G \times \dots \times G$ (n copies of G). To fix the norms, we will fix such a diffeomorphism for each n . We will mainly use the manifolds $M(= M_1), M_2$ and M_3 . To fix the norms on B (i.e. for maps from and to B), we will assume that B is a neighborhood of x_0 in a given Euclidean space. (We will shrink B whenever necessary, but the norm of the Euclidean space which contains it will not be changed).

If V_1 and V_2 are two nonnegative numbers which depend on several variables and parameters, then we will write $V_1 \preceq V_2$ (read V_1 is smaller than V_2 up to a multiplicative constant) if there is a positive constant C (which does not depend on the movable parameters and variables of V_1 and V_2) such that $V_1 \leq CV_2$. We can also write $V_1 = O(V_2)$ using Landau notation. We will write $V_1 \approx V_2$ if $V_1 \preceq V_2 \preceq V_1$.

We are interested in the C^k -topology ($k \in \mathbb{Z}_+$, $k \leq h$ if M is only C^h -smooth) of the spaces of maps from M, M_2, M_3 to $\mathfrak{g}$ and G . We will use $\|\cdot\|_k$ to denote a fixed C^k -norm. (It doesn't matter much which C^k -norm we use because they are equivalent). We use the following C^0 norms for functions from N (where N denotes one of the spaces $M, M_2, M_3, \dots$) to G :

$$(3.1) \quad \|f\|_0 = \sup_{x \in N} d(f(x), 1_G),$$

where $d(\cdot, \cdot)$ is the metric on G . To define a C^k -norm for functions from N to G , we identify TG with $\mathfrak{g} \times G$ and so on. Our convention is that the constant map from N to the neutral element in G is also denoted by 1_G ,

and $\|1_G\|_k = 0 \forall k$. Moreover, for each fixed k , if f is a map from N to $\mathfrak{g}$ with $\|f\|_0 \leq 1$ and $\exp(f)$ is the corresponding map from N to G then

$$(3.2) \quad \|f\|_k \approx \|\exp(f)\|_k .$$

Let us write down some standard inequalities which will be used later on.

If m is a fixed smooth map from N to N' (e.g., the product map from M_2 to M) and f is a map from N' to $\mathfrak{g}$ or G , then we have (for each fixed nonnegative integer k which does not exceed the smoothness class of the groupoid):

$$(3.3) \quad \|f \circ m\|_k \leq \|f\|_k .$$

If f_r from N to $\mathfrak{g}$ depends on a parameter r which lives in a probability space R , then

$$(3.4) \quad \left\| \int_R f_r dr \right\|_k \leq \sup_{r \in R} \|f_r\|_k .$$

If f_1, f_2 are two functions from N to G then we have:

$$(3.5) \quad \|f_1 \cdot f_2\|_0 \leq \|f_1\|_0 + \|f_2\|_0 ,$$

and if moreover $\|f_1\|_{k-1}, \|f_2\|_{k-1} \leq 1$ (for some fixed $k \geq 1$), then we have:

$$(3.6) \quad \|f_1 \cdot f_2\|_k \leq \|f_1\|_k + \|f_2\|_k$$

and (more refined inequalities)

$$(3.7) \quad \|f_1 \cdot f_2\|_k - \|f_2\|_k \leq \|f_1\|_k + \|f_1\|_0 \|f_2\|_k ,$$

$$(3.8) \quad \|f_1 \cdot f_2 \cdot f_1^{-1}\|_k \leq \|f_2\|_k + \|f_1\|_k \cdot \|f_2\|_0 .$$

For $f_1, f_2 : N \rightarrow \mathfrak{g}$, define $\log(\exp(f_1) \cdot \exp(f_2))$ by the Baker-Campbell-Hausdorff formula. Then we have

$$(3.9) \quad \|\log(\exp(f_1) \cdot \exp(f_2)) - f_1 - f_2\|_0 \leq \|f_1\|_0 \|f_2\|_0 ,$$

and

$$(3.10) \quad \|\exp(f_1) \cdot \exp(f_2)\|_0 \leq \|f_1 + f_2\|_0 .$$

If, moreover, $\|f_1\|_{k-1}, \|f_2\|_{k-1} \leq 1$ (for some fixed $k \geq 1$) then

$$(3.11) \quad \|\log(\exp(f_1) \cdot \exp(f_2)) - f_1 - f_2\|_k \leq \|f_1\|_0 \|f_2\|_k + \|f_2\|_0 \|f_1\|_k + \|f_1\|_{k-1} + \|f_2\|_{k-1} ,$$

and

$$(3.12) \quad \|\exp(f_1) \cdot \exp(f_2)\|_k \leq \|f_1 + f_2\|_k + \|f_1\|_0 \|f_2\|_k + \|f_2\|_0 \|f_1\|_k + \|f_1\|_{k-1} + \|f_2\|_{k-1} .$$

Finally, if f_n ($n \in \mathbb{N}$) are maps from N to G , and a_n are positive numbers such that $\sum_{n=1}^{\infty} a_n$ converges and $\|f_n\|_k \leq a_n$ (for some nonnegative integer k), then the product $f_n \cdot f_{n-1} \dots f_1$ converges in C^k -topology when $n \rightarrow \infty$ to a C^k -map from N to G .

3.2. C^0 estimates.

Lemma 3.1. *For any $\phi : M \rightarrow G$ with $\Delta(\phi) \leq 1$ we have*

$$(3.13) \quad \Delta(\widehat{\phi}) \preceq (\Delta(\phi))^2 .$$

In particular, there is a positive constant $C_0 > 0, C_0 \leq 1$ such that if $\Delta(\phi) \leq C_0$ then $\widehat{\phi}$ is well-defined and

$$(3.14) \quad \Delta(\widehat{\phi}) \leq (\Delta(\phi))^2 / C_0 \leq \Delta(\phi) .$$

Proof: Denote

$$(3.15) \quad \psi(p, q) = \phi(p.q) \cdot \phi(q)^{-1} \cdot \phi(p)^{-1} ,$$

and

$$(3.16) \quad \widehat{\psi}(p, q) = \widehat{\phi}(p.q) \cdot \widehat{\phi}(q)^{-1} \cdot \widehat{\phi}(p)^{-1} .$$

Then ψ and $\widehat{\psi}$ are functions from M_2 to G . By definition of $\widehat{\phi}$, we have

$$(3.17) \quad \begin{aligned} \widehat{\psi}(p, q) &= \exp\left(\int_{r \in T(s(q))} \log(\psi(p.q, r)) d\mu\right) \cdot \phi(p.q) \cdot \phi(q)^{-1} \cdot \\ &\quad \exp\left(\int_{r \in T(s(q))}^{-1} \log(\psi(q, r)) d\mu\right)^{-1} \cdot \phi(p)^{-1} \exp\left(\int_{r' \in T(s(p))} \log(\psi(p, r')) d\mu\right)^{-1} \\ &= \phi(p.q) \cdot \phi(q)^{-1} \cdot \phi(p)^{-1} \cdot E(p, q) = \psi(p, q) \cdot E(p, q) , \end{aligned}$$

where

$$(3.18) \quad \begin{aligned} E(p, q) &= Ad_{\psi(p,q)^{-1}} \exp\left(\int_{r \in T(s(q))} \log(\psi(p.q, r)) d\mu\right) \cdot \\ &\quad Ad_{\phi(p)} \exp\left(\int_{r \in T(s(q))}^{-1} \log(\psi(q, r)) d\mu\right)^{-1} \cdot \exp\left(\int_{r' \in T(s(p))} \log(\psi(p, r')) d\mu\right)^{-1} \\ &= \exp\left(\int_{r \in T(s(q))} \log(\psi(p, q)^{-1} \cdot \psi(p.q, r) \cdot \psi(p, q)) d\mu\right) \cdot \\ &\quad \exp\left(\int_{r \in T(s(q))} \log(\phi(p) \cdot \psi(q, r)^{-1} \cdot \phi(p)^{-1}) d\mu\right) \cdot \\ &\quad \exp\left(\int_{r \in T(s(q))} \log(\psi(p, q.r)^{-1}) d\mu\right) \quad (\text{we changed } r' \text{ by } r = q^{-1}.r') \\ &= \exp\left(\int_{r \in T} \log(A_1) d\mu\right) \cdot \exp\left(\int_{r \in T} \log(A_2) d\mu\right) \cdot \exp\left(\int_{r \in T} \log(A_3) d\mu\right) , \end{aligned}$$

where $T = T(s(q))$ and

$$(3.19) \quad \begin{aligned} A_1 &= \psi(p, q)^{-1} \cdot \psi(p.q, r) \cdot \psi(p, q) , \\ A_2 &= \phi(p) \cdot \psi(q, r)^{-1} \cdot \phi(p)^{-1} , \\ A_3 &= \psi(p, q.r)^{-1} . \end{aligned}$$

One verifies directly that

$$(3.20) \quad A_1 \cdot A_2 \cdot A_3 = \psi(p.q)^{-1} .$$

Consider A_1, A_2, A_3 as maps from M_3 to G . By definition, $\Delta(\phi) = \|\psi\|_0$. The inequality $\Delta(\phi) \leq 1$ in the hypothesis of Lemma 3.1, together with the fact that the metric on G is bi-invariant, implies that

$$(3.21) \quad \|A_1\|_0 = \|A_2\|_0 = \|A_3\|_0 = \|\psi\|_0 = \Delta(\phi) \leq 1$$

Applying Inequalities (3.9), (3.4) and (3.21) several times to $E(p, q)$, we get:

$$(3.22) \quad \begin{aligned} \log E(p, q) &= \varepsilon_1 + \int_{r \in T} \log(A_1) d\mu + \int_{r \in T} \log(A_2) d\mu + \int_{r \in T} \log(A_3) d\mu \\ &= \varepsilon_1 + \int_{r \in T} [\log(A_1) + \log(A_2) + \log(A_3)] d\mu \\ &= \varepsilon_1 + \varepsilon_2 + \int_{r \in T} \log(A_1 A_2 A_3) d\mu \\ &= \varepsilon_1 + \varepsilon_2 - \log(\psi(p, q)) \end{aligned}$$

where ε_1 and ε_2 are some functions such that

$$(3.23) \quad \|\varepsilon_1\|_0, \|\varepsilon_2\|_0 \preceq \Delta(\phi)^2$$

In other words, we have $\|\log(\psi(p, q)) + \log E(p, q)\|_0 \preceq \Delta(\phi)^2$, which implies, by Inequality (3.10), that $\|\psi.E\|_0 \preceq \Delta(\phi)^2$. But we have $\widehat{\psi}(p, q) = \psi(p, q).E(p, q)$, therefore

$$(3.24) \quad \Delta(\widehat{\phi}) = \|\widehat{\psi}\|_0 = \|\psi.E\|_0 \preceq \Delta(\phi)^2$$

□

Lemma 3.1 immediately implies the uniform convergence (i.e. convergence in C^0 topology) of the sequence of maps $\phi_n : M \rightarrow G$, defined iteratively by $\phi_{n+1} = \widehat{\phi_n}$, beginning with an arbitrary smooth map ϕ_1 which satisfies the inequality $\Delta(\phi_1) \leq C_0/4$. (This inequality can always be achieved by replacing B by a sufficiently small invariant neighborhood of x_0 in B if necessary). Indeed, since $\|\psi_2\|_0 = \Delta(\phi_2) \leq (\Delta(\phi_1))^2/C_0 \leq \Delta(\phi_1) \leq C_0/4 \leq 1/4$ by Lemma 3.1, where

$$(3.25) \quad \psi_n(p, q) = \phi_n(p.q) \cdot \phi_n(q)^{-1} \cdot \phi_n(p)^{-1},$$

we can define $\phi_2 = \widehat{\phi_1}$, and so on, hence ϕ_n is well defined for all $n \in \mathbb{N}$. By recurrence on n , one can show easily that we have

$$(3.26) \quad \|\psi_n\|_0 \leq C_0 \cdot (b_0)^{2^n} \quad \forall n \in \mathbb{N}, \quad \text{where } b_0 = \frac{1}{2} < 1,$$

which implies in particular that $\sum_{n=1}^{\infty} \|\psi_n\|_0 < \infty$ (this is a very fast converging series). Put

$$(3.27) \quad \Psi_n(p, q) = \exp\left(\int_{q \in T(s(p))} \log(\psi_n(p.q)) d\mu\right).$$

Then $\|\Psi_n\|_0 \preceq \|\psi_n\|_0$ (by Inequalities (3.3) and (3.4)), which together with $\sum_{n=1}^{\infty} \|\psi_n\|_0 < \infty$ implies that

$$(3.28) \quad \sum_{n=1}^{\infty} \|\Psi_n\|_0 < \infty .$$

This last inequality implies the convergence of the product $\Psi_n \cdot \Psi_{n-1} \dots \Psi_1$ in C^0 -topology when $n \rightarrow \infty$. But

$$(3.29) \quad \Psi_n \cdot \Psi_{n-1} \dots \Psi_1 = \phi_{n+1} \cdot \phi_1^{-1}$$

Thus ϕ_n converges in C^0 -topology when $n \rightarrow \infty$. Denote by ϕ_{∞} the limit

$$(3.30) \quad \phi_{\infty} = \lim_{n \rightarrow \infty} \phi_n$$

Then ϕ_{∞} is a continuous homomorphism from M to G . It is also clear that the restriction of ϕ_{∞} to G is the identity map from G to itself.

It remains to show that ϕ_{∞} is smooth. This is the purpose of the next subsection, where we will show that for any $k \in \mathbb{N}$, $k \leq h$ if M belongs to the class C^h only, we have $\phi_{\infty} = \lim_{n \rightarrow \infty} \phi_n$ in C^k -topology as well.

Remark 3.2. When G is Abelian then the above computations show that $\widehat{\psi} = 1_G$, i.e. $\widehat{\phi}$ is already a homomorphism from M to G , and there is no need to iterate our averaging process.

3.3. C^k estimates. Roughly speaking, we want to make estimations on ψ_n in order to show that, if k does not exceed the smoothness class of the groupoid $M \rightrightarrows B$, then $\sum_{n=1}^{\infty} \|\psi_n\|_k < \infty$. If this series converges, then similarly to the previous subsection, we also have $\sum_{n=1}^{\infty} \|\Psi_n\|_k < \infty$ where $\Psi_n = \phi_{n+1} \cdot \phi_n^{-1}$ is given by formula (3.27), hence the product $\Psi_n \cdot \Psi_{n-1} \dots \Psi_1$ converges in C^k -topology when $n \rightarrow \infty$, implying that $\phi_n \rightarrow \phi_{\infty}$ in C^k -topology.

Lemma 3.3. *Let $k \in \mathbb{N}$ be a natural number which does not exceed the smoothness class of the groupoid $M \rightrightarrows B$. Assume that $\|\psi\|_0 = \Delta(\phi) \leq 1$ and $\|\phi\|_{k-1} \preceq 1$. Then we have:*

$$(3.31) \quad \|\widehat{\psi}\|_k \preceq \|\psi\|_0 \|\psi\|_k + \|\psi\|_{k-1} + \|\psi\|_0 \|\phi\|_{k-1} + \|\psi\|_0^2 \|\phi\|_k$$

Proof. Assume that $\|\phi\|_{k-1} \preceq 1$ by hypothesis of Lemma 3.3. Then by Inequality (3.6) and Inequality (3.3), we have $\|\psi\|_{k-1} \preceq \|\phi\|_{k-1} \preceq 1$. Let A_1, A_2, A_3 be the functions defined by Equation (3.19). We want to estimate them. For $A_3 = \psi(p, q.r^{-1})^{-1}$, using Inequality (3.3), we get:

$$(3.32) \quad \|A_3\|_{k-1} \preceq \|\psi\|_{k-1} \preceq 1 \quad \text{and} \quad \|A_3\|_k \preceq \|\psi\|_k .$$

For $A_1 = \psi(p, q)^{-1} \cdot \psi(p, q, r) \cdot \psi(p, q)$, using Inequality (3.6) (and Inequality (3.3)), we also get

$$(3.33) \quad \|A_1\|_{k-1} \preceq \|\psi\|_{k-1} \preceq 1 \quad \text{and} \quad \|A_1\|_k \preceq \|\psi\|_k.$$

The estimation of $A_2 = \phi(p) \cdot \psi(q, r)^{-1} \cdot \phi(p)^{-1}$ is more complicated, because it involves the function ϕ directly. Using Inequality (3.8) we get

$$(3.34) \quad \|A_2\|_{k-1} \preceq \|\psi\|_{k-1} + \|\phi\|_{k-1} \|\psi\|_0 \preceq 1$$

and

$$(3.35) \quad \|A_2\|_k \preceq \|\psi\|_k + \|\phi\|_k \|\psi\|_0.$$

Applying Inequality (3.11) and the above inequalities to $E(p, q)$, we get that

$$(3.36) \quad \|\varepsilon_1\|_k, \|\varepsilon_2\|_k \preceq (\|\psi\|_k + \|\phi\|_k \|\psi\|_0) \|\psi\|_0 + (\|\psi\|_{k-1} + \|\phi\|_{k-1} \|\psi\|_0)$$

Moreover, we have

$$(3.37) \quad \|E\|_i \preceq \|\psi\|_i + \|\psi\|_0 \|\phi\|_i \quad \forall i = 0, \dots, k.$$

Now applying Inequality (3.12) and the last two inequalities, we get

$$(3.38) \quad \begin{aligned} \|\widehat{\psi}\|_k &= \|\exp(\log(\psi)) \cdot \exp(\log(E))\|_k \\ &\preceq \|\log(\psi) + \log(E)\|_k + \|\psi\|_0 \|E\|_k + \|\psi\|_k \|E\|_0 + \|\psi\|_{k-1} + \|E\|_{k-1} \\ &= \|\varepsilon_1 + \varepsilon_2\|_k + \|\psi\|_0 \|E\|_k + \|\psi\|_k \|E\|_0 + \|\psi\|_{k-1} + \|E\|_{k-1} \\ &\preceq \|\psi\|_0 \|\psi\|_k + \|\psi\|_{k-1} + \|\psi\|_0 \|\phi\|_{k-1} + \|\psi\|_0^2 \|\phi\|_k \end{aligned}$$

□

Lemma 3.4. *With the assumptions of Lemma 3.3 we have:*

$$(3.39) \quad \|\widehat{\phi}\|_k - \|\phi\|_k \preceq \|\psi\|_k + \|\psi\|_0 \|\phi\|_k$$

Proof. Applying Inequality (3.7) to $\widehat{\phi} = \Psi \cdot \phi$, we get

$$(3.40) \quad \|\widehat{\phi}\|_k - \|\phi\|_k \preceq \|\Psi\|_k + \|\Psi\|_0 \|\phi\|_k$$

Now replace $\|\Psi\|_0$ by $\|\psi\|_0$ and $\|\Psi\|_k$ by $\|\psi\|_k$. □

Lemma 3.5. *Assume that ϕ_1 is a map from M to G such that $\Delta(\phi_1) < C_0/4$, and that $\phi_{n+1} = \widehat{\phi}_n$ for any $n \in \mathbb{N}$, as in the previous subsection. Let k be a natural number which does not exceed the smoothness class of the groupoid $M \rightrightarrows B$. Then there is a finite positive number $D_k > 0$ and a positive number $0 < b_k < 1$, such that for any $n \in \mathbb{N}$ the following two inequalities hold:*

$$(3.41) \quad \|\phi_n\|_k \leq D_k \cdot (1 - 2^{-n})$$

and

$$(3.42) \quad \|\psi_n\|_k \leq D_k (b_k)^{2^n}.$$

Proof of Theorem 1.1. Remark that if Inequality (3.41) is satisfied, then in particular the k -norms $\|\phi_n\|_k$ ($n \in \mathbb{N}$) is bounded by D_k , and this is enough to imply (by Ascoli theorem) that $\phi_\infty = \lim_{n \rightarrow \infty} \phi_n$ is of class C^k . Inequality (3.42) is also a sufficient condition for the C^k -smoothness of ϕ_∞ , because it implies in particular that $\sum_{n=1}^{\infty} \|\Psi_n\|_k \preceq \sum_{n=1}^{\infty} \|\psi_n\|_k < \infty$, which in turns implies that the sequence of maps (ϕ_n) converges in C^k -topology. Thus the smoothness of the homomorphism $\phi_\infty : M \rightarrow G$, and hence Theorem 1.1, follows directly from the above Lemma, and in fact we need only one of the above two inequalities (3.42) and (3.41) in order to prove our main result. But these two inequalities are closely related (they need each other in a proof by induction), so we put them together. $\square$

Proof of Lemma 3.5. We will prove it by induction on k . When $k = 0$, Lemma 3.5 is already proved in the previous section (with $b_0 = 1/2$). Let us now assume that Inequalities (3.42) and (3.41) are true at the level $k - 1$ (i.e. if we replace k by $k - 1$). We will show that they are true at the level k .

We will choose an (arbitrary) number $b_k > 0$ such that $1 > b_k > b_k^2 > b_{k-1}, b_0$. (For example, one can put $b_0 = 1/2$ and then $b_k = (b_{k-1})^{1/3}$ by recurrence). What will be important for us is that $b_0/b_k, b_{k-1}/b_k^2$ and b_0/b_k^2 are positive numbers which are strictly smaller than 1.

It follows from Lemma 3.3 and Lemma 3.4 that there exist two positive numbers c_1 and c_2 (which do not depend on n) such that we have, for any $n \in \mathbb{N}$:

$$(3.43) \quad \|\psi_{n+1}\|_k \leq c_1 (\|\psi_n\|_0 \|\psi_n\|_k + \|\psi_n\|_{k-1} + \|\psi_n\|_0 \|\phi_n\|_{k-1} + \|\psi_n\|_0^2 \|\phi_n\|_k)$$

and

$$(3.44) \quad \|\phi_{n+1}\|_k - \|\phi_n\|_k \leq c_2 (\|\psi_n\|_k + \|\psi_n\|_0 \|\phi_n\|_k).$$

We will now prove Inequalities (3.42) and (3.41) by induction on n . There exists a natural number n_0 such that for any $n > 0$ we have

$$(3.45) \quad Q_1 := D_0 \left(\frac{b_0}{b_k} \right)^{2^n} + \left(\frac{b_{k-1}}{b_k^2} \right)^{2^n} + D_0 \left(\frac{b_0}{b_k^2} \right)^{2^n} + D_0^2 \left(\frac{b_0}{b_k} \right)^{2^{n+1}} \leq \frac{1}{c_1}$$

and

$$(3.46) \quad Q_2 := (b_k)^{2^n} + D_0 (b_0)^{2^n} \leq \frac{2^{-n-1}}{c_2}.$$

By choosing D_k large enough, we can assume that Inequalities (3.42) and (3.41) are satisfied for any $n \leq n_0$. We will also assume that $D_k \geq D_{k-1}$. let us now show that if Inequalities (3.42) and (3.41) are satisfied for some $n \geq n_0$ then they are still satisfied when we replace n by $n + 1$. (This is the last step in our induction process).

Indeed, for $\|\psi_{n+1}\|_k$, using Inequality (3.43) and the induction hypothesis, we get

$$\begin{aligned}
 (3.47) \quad \|\psi_{n+1}\|_k &\leq c_1 (\|\psi_n\|_k \|\psi_n\|_0 + \|\psi_n\|_{k-1} + \|\psi_n\|_0 \|\phi_n\|_{k-1} + \|\psi_n\|_0^2 \|\phi_n\|_k) \\
 &\leq c_1 \left(D_k(b_k)^{2^n} D_0(b_0)^{2^n} + D_{k-1}(b_{k-1})^{2^n} + D_0(b_0)^{2^n} D_{k-1} + D_0^2(b_0)^{2^{n+1}} D_k \right) \\
 &\leq D_k c_1 \left((b_k)^{2^n} D_0(b_0)^{2^n} + (b_{k-1})^{2^n} + D_0(b_0)^{2^n} + D_0^2(b_0)^{2^{n+1}} \right) \\
 &= D_k c_1 Q_1(b_k)^{2^{n+1}} \\
 &\leq D_k (b_k)^{2^{n+1}}
 \end{aligned}$$

Similarly, for $\|\phi_{n+1}\|_k$ we have:

$$\begin{aligned}
 (3.48) \quad \|\phi_{n+1}\|_k &\leq \|\phi_n\|_k + c_2 (\|\psi_n\|_k + \|\psi_n\|_0 \|\phi_n\|_k) \\
 &\leq D_k(1 - 2^{-n}) + c_2 (D_k(b_k)^{2^n} + D_0(b_0)^{2^n} \cdot D_k) \\
 &\leq D_k(1 - 2^{-n}) + D_k c_2 [(b_k)^{2^n} + D_0(b_0)^{2^n}] \\
 &= D_k(1 - 2^{-n}) + D_k c_2 Q_2 \\
 &\leq D_k(1 - 2^{-n}) + D_k 2^{-n-1} = D_k(1 - 2^{-n-1})
 \end{aligned}$$

□

Remark 3.6. If we start with a near-homomorphism from M to a compact Lie group H different from G , then our iterative averaging method still yields a homomorphism from M to H . Thus, in a sense, our result generalizes a result of Grove, Karcher and Ruh [6] on near-homomorphisms between compact groups.

Acknowledgements. I would like to thank Alan Weinstein for getting my attention to the problem, and for many stimulating discussions. Thanks to Philippe Monnier for finding some errors in the first version of this paper.

REFERENCES

1. C. Anantharaman-Delaroche, J. Renault, Amenable groupoids. Monographies de l'Enseignement Mathématique, 36 (2000).
2. Salomon Bochner, *Compact groups of differentiable transformations*, Annals of Math. (2) **46** (1945), 372–381.
3. Elie Cartan, *Leçons sur la géométrie des espaces de Riemann*, Gauthiers-Villars, Paris, 1928.
4. ———, *Groupes simples clos et ouverts et géométrie riemannienne*, J. Math. Pures Appl. **8** (1929), 1–33.
5. M. Crainic and R. Fernandes, *Integrability of Lie brackets*, preprint math.DG/0105033, Annals of Math. (to appear).
6. K. Grove, H. Karcher, and E. Ruh, *Group actions and curvature*, Invent. Math. **23** (1974), 31–48.
7. Ph. Monnier and Nguyen Tien Zung, *Levi decomposition for smooth Poisson structures*, preprint math.DG/0209004.
8. Alan Weinstein, *Almost invariant submanifolds for compact group actions.*, J. Eur. Math. Soc. (JEMS) **2** (2000), no. 1, 53–86.

9. ———, *Linearization problems for Lie algebroids and Lie groupoids*, Lett. Math. Phys. **52** (2000), no. 1, 93–102.
10. ———, *Poisson geometry of discrete series orbits, and momentum convexity for non-compact group actions*, Lett. Math. Phys. **56** (2001), No.1, 17–30.
11. ———, *Linearization of regular proper groupoids*, J. Inst. Math. Jussieu **1** (2002), no. 3, 493–511.
12. Nguyen Tien Zung, *Levi decomposition of analytic Poisson structures and Lie algebroids*, Topology **42** (2003), No. 6, 1403–1420.
13. ———, *Torus actions and integrable systems*, preprint 2003
14. ———, *Proper groupoids and Poisson convexity*, in preparation.

LABO. EMILE PICARD, UMR 5580 CNRS, UFR MIG, UNIVERSITÉ TOULOUSE III
 E-mail address: `tienzung@picard.ups-tlse.fr`